



Effects of phosphate on growth and skeletal density in the scleractinian coral *Acropora muricata*: A controlled experimental approach

Jeremy G. Dunn ^{a,b,c,1}, Paul W. Sammarco ^{a,c,*}, Gary LaFleur Jr. ^a

^a Department of Biological Sciences, Nicholls State University, Thibodaux, LA 70310, USA

^b Coastal Estuary Services, LLC, Shaw Environmental & Infrastructure Group, 11027 LA Hwy. 35, Kaplan, LA 70548, USA

^c Louisiana Universities Marine Consortium (LUMCON), 8124 Hwy. 56, Houma, LA 70344, USA

ARTICLE INFO

Article history:

Received 30 April 2011

Received in revised form 3 October 2011

Accepted 4 October 2011

Available online 24 November 2011

Keywords:

Acropora muricata

Coral

Density

Growth

Phosphate

ABSTRACT

Phosphate contamination can negatively affect corals, modifying growth rates, skeletal density, reproduction, mortality, and zooxanthellae. We determined the effects of elevated phosphate on coral growth and density. Genetically distinct colonies of *Acropora muricata* were sub-divided and distributed among three 110-L aquaria, and exposed to phosphate levels of 0.09, 0.20, and 0.50 mg L⁻¹ for four months. Total skeletal length, living tissue length, weight, branch production, and polyp extension were measured. Linear extension and tissue growth increased under all conditions. Growth rates were highest at a phosphate concentration of 0.50 mg L⁻¹. Weight increased through time, graded from low to high with phosphate concentration. Density decreased through time, and was significantly lowest in the high phosphate treatment. Phosphate concentration produced no visible effects of stress on the corals, as indicated by polyp extension and lack of mortality. It is suggested that the phosphate enhanced growth was due to increased zooxanthellar populations and photosynthetic production within the coral. Skeletal density reduction may be due to phosphate binding at the calcifying surface and the creation of a porous and structurally weaker calcium carbonate/calcium phosphate skeleton. Increased phosphate concentrations, often characteristic of eutrophic conditions, caused increased coral growth but also a more brittle skeleton. The latter is likely more susceptible to breakage and damage from other destructive forces (e.g., bioerosion) and makes increased coral growth a poor indicator of reef health.

© 2011 Published by Elsevier B.V.

1. Introduction

1.1. Background

Coral reefs are one of the most productive and biodiverse ecosystems in the world (Jackson, 1991; Moberg and Folke, 1999; Reaka-Kudla, 1997; Roberts et al., 2002; Sebens, 1994). Unfortunately, coral colonies are disappearing at an alarming rate. Ten percent of the world's reefs are already seriously damaged (Miller, 1999; also see Knowlton and Jackson, 2008; Tamelander and Rajasuriya, 2008; Schutte et al., 2010), and sixty percent are predicted to exhibit serious damage within the next 20–40 years. Causes of this damage include increased sea surface temperatures (Baker et al., 2008; Hoegh-Guldberg, 1999; Wild et al., 2010), disease (Aeby et al., 2011; Aronson and Precht, 2001; Green and Bruckner, 2000; Patterson et al., 2002; Williams and Miller, 2005), overfishing of reef fish (Hay and Rasher, 2010; McManus 1997), mortality of grazing sea urchins in the Caribbean (Hughes, 1994; Hughes et al., 2010; Sammarco 1980, 1982a, 1982b),

urban development (Finkl and Charlier, 2003; Hawkins and Roberts, 1994; Reopanichkul et al., 2010), exploitation for the aquarium trade (Laurance, 2010; Wood, 2001), and physical damage from divers (Guzner et al., 2010; Juhasz et al., 2010; Zakai and Chadwick-Furman, 2002).

Eutrophication can also cause coral mortality and sub-lethal effects on coral reefs (Hunter and Evans, 1995; Hunter and Stephenson, 1999; Lapointe et al., 1990, 2004; Rothenberger et al., 2008; Smith et al., 1981). Coastal waters can become eutrophic due to the addition of fertilizer (DeGeorges et al., 2010; Fishelson, 1973), sewage (Costa et al., 2000; DeGeorges et al., 2010), and other terrestrial run-off (Fabricius, 2005), increasing phosphorus and nitrogen concentrations in nearby reef systems (Costa et al., 2000; Lapointe et al., 2003, 2010). Human populations are increasing along coastal regions, with about two-thirds of the world's population (~3.9 billion people) now residing within 160 km of the coast. This number is estimated to increase to 6.2 billion by 2025 (Miller, 1999).

Corals possess endosymbiotic zooxanthellae (genus *Symbiodinium*) which receive metabolic wastes in the form of phosphorus, nitrogen, and carbon dioxide from the coral, while the coral receives oxygen and carbohydrates from the photosynthetic activities of the zooxanthellae (Baker, 2003; Barnes and Hughes, 1982; Furla et al., 2011; Trench, 1993; Yonge et al., 1932). The symbiotic relationship between coral and zooxanthellae has been well studied, although the complete details

* Corresponding author at: Louisiana Universities Marine Consortium (LUMCON), 8124 Hwy. 56, Houma, LA 70344, USA. Tel.: +1 985 851 2876; fax: +1 985 851 2874.
E-mail address: psammarco@lumcon.edu (P.W. Sammarco).

¹ Current address.

of this interaction are still unclear. The coral hosts are clearly dependent upon the zooxanthellae for survival, because the zooxanthellae transfer up to 99% of their photosynthetic products to the coral via translocation (Muscattine and Cernichiar, 1969;) of such products as glycerol, glucose, and alanine (Barnes, 1987; Falkowski et al., 1984; Muscattine and Cernichiar, 1969; Stambler, 2011). Although corals obtain up to 95% of their carbon requirements from the zooxanthellae (Muscattine, 1990; Muscattine et al., 1981), many species still must acquire additional nutrients and energy from zooplankton (Johannes et al., 1970). The degree of dependence upon zooxanthellar products varies according to coral species (Porter, 1976) and depth of the colony (Falkowski et al., 1984; Muscattine et al., 1989). For example, massive corals (e.g. *Montastraea*) can be more heterotrophic, while some branching corals (e.g. *Acropora*) rely mainly on zooxanthellae to meet their energy needs (Porter, 1976). Zooxanthellar photosynthesis and coral calcification are tightly linked (Al-Horani et al., 2003; Allemand et al., 2004; McConnaughey and Whelan, 1997; Suzuki et al., 1995).

Zooxanthellae require nutrients (P and N), dissolved inorganic carbon (DIC), and solar radiation for photosynthesis. Hermatypic corals require, minimally, DIC, ATP, and calcium for calcification.

The endosymbiotic zooxanthellae are concentrated in the oral endodermal cells surrounding the coelenteron of the coral polyp (Allemand et al., 2004). Changes in rates of calcification, photosynthesis, exchange of seawater through the mouth, and trans-epithelial ionic transport can affect the chemistry of the coelenteron (Gattuso et al., 1999). Expansion or retraction of coral polyps can affect calcium transport via the mouth. Calcium ions enter into the coelenteron via the mouth and/or by transport across the oral epithelium. Elevated phosphate levels in seawater may disrupt normal carbon cycle functions and concentrations of compounds produced and shared within the holobiont, resulting in abnormal growth. The chemistry of the coelenteron can differ from that of the surrounding seawater due to internal processes (e.g. photosynthesis, respiration, calcification; Al-Horani et al., 2003), as well as reduced flushing of water through the mouth (Wright and Marshall, 1991). Recently, it was determined that P/Ca ratios in coral skeletons provide a record of phosphate concentrations and thus nutrient availability in the water through geological time (LaVigne et al., 2008, 2010).

1.2. Effects of phosphate enrichment on corals

Average phosphorus concentrations on natural reefs are generally low (0.01 mg L^{-1} ; Kleypas et al., 1999), but there appears to be a trend towards increasing values in recent years. For example, concentrations up to 0.266 mg L^{-1} have been documented on some reefs in the Southern Gulf of Mexico (Cruz-Piñón et al., 2003). Elevated phosphate levels are known to affect several aspects of coral health, such as growth rate (Bucher and Harrison, 2002; Ferrier-Pagès et al., 2000; Koop et al., 2001; Renegar and Riegl, 2005; Steven and Broadbent, 1997), skeletal density (Koop et al., 2001; Rasmussen et al., 1993), reproduction (Harrison and Ward, 2001; Koop et al., 2001), mortality (Koop et al., 2001; Walker and Ormond, 1982), disease susceptibility (Bruno et al., 2003), and zooxanthellar density (Koop et al., 2001; Steven and Broadbent, 1997; Takabayashi, 1996). Phosphate can stimulate phytoplankton population growth, potentially reducing the availability of light to corals (Walker and Ormond, 1982), or stimulating growth of benthic filamentous algae that can overgrow corals (Done, 1992). Overgrowth by filamentous algae can be particularly detrimental in areas where populations of grazers such as fish and sea urchins have been dramatically reduced by overfishing or disease (Hughes and Connell, 1999).

The effects of phosphate on coral condition and/or growth have been studied both in the lab and the field (Ferrier-Pagès et al., 2000; Kinsey and Davies, 1979; Koop et al., 2001; Rasmussen, 1986, 1988; Rasmussen et al., 1993; Renegar and Riegl, 2005; Stambler et al., 1991). The results of past studies, however, were not consistent, particularly with respect to coral growth. Growth is often considered an

indicator of coral health (Neudecker, 1983; Rosen, 1981) and is commonly quantified via measurement of linear extension (Jinendradasa and Ekaratne, 2002) or changes in buoyant weight (Davies, 1989). Although the buoyant weight technique is attractive because it allows for non-destructive measurement of complete colony growth, it was not used in our study, because it may produce highly variable data (Bucher et al., 1998).

Koop et al. (2001) performed a study monitoring effects of nutrient enrichment on coral reefs in the field (called Effect of Nutrient Enrichment on Coral Reefs Experiment – ENCORE), where phosphate levels were manipulated over two one-year periods on the Great Barrier Reef. In the first year, low doses of phosphate (0.218 mg L^{-1} , declining to $0.0475 \text{ mg L}^{-1} \text{ PO}_4^{3-}$ within 2–3 h) were introduced to a semi-enclosed “micro-atoll”. In the second year, higher phosphate concentrations were used (0.484 mg L^{-1} declining to $0.228 \text{ mg L}^{-1} \text{ PO}_4^{3-}$ at the end of low tide). Elevated phosphate concentrations did affect coral health, but these effects varied between coral species and experimental phosphate levels. For example, linear extension rates increased to 49 mm/yr in *Acropora longicyathus* at a phosphate concentration of 0.484 mg L^{-1} , but not at a lower concentration of 0.218 mg L^{-1} (Bucher and Harrison, 2002; Koop et al., 2001). No increases in growth were observed in the other coral species considered (Koop et al., 2001; Takabayashi, 1996). Similarly, Stambler et al. (1991) used phosphorus concentrations similar to Koop et al.’s (2001; 0.009 mg L^{-1} , 0.048 mg L^{-1} , 0.190 mg L^{-1}) and found no changes in linear extension rate in *Pocillopora damicornis*.

Bell et al. (2007) re-considered Koop et al.’s (2001) work and found that waters associated with their experimental reefs were already above the defined target experimental values, placing the results of this experiment in question. In considering algal growth, Bell et al. felt that this was the reason that there was no significant response in algal growth observed to the nutrient additions (they note anecdotally, however, that algal growth increased on the crests and sides of these reefs, indicating that nutrient enrichment does indeed affect algal growth).

In past studies, a buoyancy technique (Davies, 1989) was used to estimate calcification rates in *Acropora longicyathus* (Bucher and Harrison, 2002; Koop et al., 2001) and *A. palifera* (Koop et al., 2001; Steven and Broadbent, 1997). Under the highest phosphorus concentrations, both of these species exhibited increased calcification rates. *A. aspera* did not (Koop et al., 2001), indicating a species-specific response. More variability was seen when calcification rates were documented to decrease in *Stylophora pistillata* and *P. damicornis* exposed to high phosphorus concentrations (Koop et al., 2001; Takabayashi, 1996). Furthermore Ferrier-Pagès et al. (2000) found that fragments of *S. pistillata* exhibited a 60% decrease in estimated calcification rates at phosphorus concentrations of 0.266 mg L^{-1} . Similarly, Renegar and Riegl (2005) observed decreased rates of calcification in *Acropora cervicornis* exposed to phosphate concentrations of 0.393 mg L^{-1} , with no effect at 0.204 mg L^{-1} . Exposure to increased phosphorus concentrations caused a significant reduction in both skeletal density (i.e., greater porosity) and calcification rates in *A. longicyathus* (Bucher and Harrison, 2002; Koop et al., 2001). On the other hand, the same concentrations caused significant increases in density in other coral species (Koop et al., 2001; Takabayashi, 1996).

In a field study, Risk and Sammarco (1991) reported that the coral *Porites lobata* exhibited increasing density with distance from shore on the Great Barrier Reef. Inshore corals were characterized by a lower skeletal density than those offshore. They also demonstrated a higher skeletal porosity inshore. Scanning electron microscopy (SEM) examination of skeletal micro-architecture revealed that the difference in coral density was not due to a decrease in the number of septa and dissepiments, or a change in the spacing between them; rather, it was due to the differences in porosity and the thickness of the calical characters. Risk and Sammarco (1991) suggested that this inshore–offshore cline could have been caused by higher

levels of inshore nutrients, primarily phosphate which has been suspected to hinder calcification (Kinsey and Davies, 1979; Sammarco and Risk, 1990).

In the present study, we utilized a set of laboratory microcosms to examine the effects of phosphate on corals. We controlled heavily for a variety of potential confounding factors, including phytoplankton populations (removed by filtration), filamentous algae (removed with grazing snails), light – diurnal and nocturnal (controlled with precise timing and monitoring of radiant exposure), temperature (maintained at a constant of 25 °C), PO_4 concentration (maintained with daily monitoring and adjustment), pH (maintained at 8.2), O_2 concentration (maintained at saturated level), salinity (maintained at 35 ppt with an automatic top-up system), and nitrogenous compounds (kept in check using a bacterial filter). In addition, we employed regular and frequent water quality sampling with a high level of accuracy and precision, large tanks (to decrease micro-scale variance), a small number of fragments per tank (to decrease crowding and fouling effects), frequent feeding of corals (to reduce potential suppressed growth rates common in captivity), genetic diversity of colonies (confirmed using DNA analysis), using weight and total skeletal linear length to calculate density, and running the experiment for an extensive period of time – ~4 months.

Using this controlled experimental approach, we attempted to determine the effects of increased phosphate concentration in seawater on the hermatypic zooxanthellate scleractinian coral *Acropora muricata* (see Wallace, 1999 for discussion on the taxonomy for this species formerly known as *Acropora formosa*) in a controlled laboratory environment. We did this by adding phosphate to aquarium microcosms at three different levels and monitoring total skeletal length, living tissue length, weight, number of branches, and degree of polyp extension. Through analyses of these variables, we demonstrate that, although skeletal growth and weight increase in these corals under high phosphate conditions, skeletal density decreases, potentially compromising skeletal integrity. We also include a discussion of possible physiological mechanisms driving this response.

2. Materials and methods

2.1. Target species

The coral species used in this experiment was the branching coral *Acropora muricata*, one of the most rapidly growing stony corals

recorded, with an average annual linear extension rate of up to 121 mm/yr (Jinendradasa and Ekaratne, 2002). This Indo-Pacific species can be readily purchased. The five colonies used here came from Fiji and were obtained from Live Aquaria (Rhineland, WI).

2.2. Experimental design

The experiment followed a mixed model, repeated measures, replicated, unbalanced orthogonal design. It was run for 115 days and consisted of exposing fragments of *Acropora muricata* to varying phosphate concentrations (0.09, 0.20, 0.50 mg L^{-1}) – concentrations similar to those used in previous studies (Ferrier-Pagès et al., 2000; Koop et al., 2001; Renegar and Riegl, 2005). Five coral colonies were fragmented into 27 sub-colonies (Fig. 1). The resulting ramets (Futuyma, 1998) were randomly distributed among three 110 L treatment tanks (Fig. 2).

2.3. Procedure and processing of coral

We confirmed through molecular genetic analysis using Amplified Fragment Length Polymorphisms (AFLPs) that our colonies were all genetically distinct. Samples were processed at the Dept. of Pharmaceutical Sciences, Univ of Buffalo, NY using the protocols described in Atchison (2005), Brazeau et al. (2005) and Atchison et al. (2008).

Live corals were maintained in the Climate Control Laboratory, Department of Biological Sciences, Nicholls State University (Fig. 2). Coral colonies were fragmented into ramets using bone cutters, secured onto retractable PVC piping via epoxy putty (AquaStick®), and distributed randomly into separate treatment tanks attached to a PVC support-base and labeled using a soldering iron (Fig. 3).

2.4. Water chemistry

Phosphate levels (mg L^{-1}) in the seawater were measured three times per week (average) to maintain target concentrations. Ammonia (N; mg L^{-1}), nitrite (NO_2 ; mg L^{-1}), pH, temperature (°C), and salinity (ppt) were recorded at these times. Nitrate-N (mg L^{-1}) and calcium (mg L^{-1}) levels were measured at weekly intervals (average). Phosphate levels were measured using Hach® Phos Ver3 reagents (Cat. 2125-99). In cases where phosphate concentrations were low, phosphate was added and re-tested after 30 min. Phosphate levels were adjusted using sodium phosphate ($\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$;




Fig. 1. Schematic of experimental design. Five genetically distinct coral colonies were fragmented and distributed into three tanks holding varying experimental phosphate levels.




Fig. 2. Experimental tank showing PVC support structure with coral fragments. Tank fitted with three powerheads, filter attachment, protein skimmer and live rock/algal nitrogen control system (from Fiji; upper right), heaters, salinity stabilization system, and UV sterilizer. Lighting provided via two 400 W metal halide bulbs and two 160 W VHO actinic bulbs and cooling fans. Timers simulated a natural 12:12 h, light: dark photoperiod, duration – 5 h of high intensity metal halide lighting and 12 h of actinic lighting per day.




Fig. 3. (A) PVC insert with coral fragment. Measurements taken from the coral included total skeletal linear length (TSL; mm), living linear length (LL; mm), weight (g), number of branches, and polyp extension. (B) PVC support structure for corals, with holders inserted.

Acros Organics®, NJ, Product # 42440). pH levels were maintained between 8.2 and 8.4, using a synthetic saltwater buffer (Reef Buffer, Seachem®). Ammonia and nitrite levels were measured using Jungle Labs® test kits, and Hach reagents (Nitra®Ver 5) were used to measure nitrate levels. (Calcium levels were measured using test kits by Salifert and supplemented using a calcium additive; Reef Complete, Seachem®.) Saltwater (Instant Ocean®) was prepared with water from a reverse-osmosis deionization unit (RODI; Air Water Ice®, Extreme Typhoon 150 III), and salinity was maintained between 34 and 35 ppt, being measured with a handheld refractometer.

2.5. Equipment

A. muricata colonies require a high degree of lighting, strong turbulent water flow, high calcium concentrations, and a steady temperature (Borneman, 2001). Water flow was provided by two powerheads (Maxi-Jet® 1200) and powered by a wave-making unit (Maxi-Jet® Natural Wavemaker System). Powerheads were timed to turn on and off at 20-s intervals, creating a pulsing flow of current. Magnetic powerhead holders (Algae Free® Sure Grip 50) were used to secure the powerheads to the glass. Powerheads (Hagen® Aqua Clear 50) fitted with a quick filter attachment provided additional water flow debris filtering. Protein skimmers (CPR® Bak-Pak 2R) removed excess waste accumulation, particularly organic matter. Heaters (Marineland® Visi-Therm Stealth 250 W) were used to maintain temperatures at 25–26 °C. Tanks were covered with glass to reduce evaporation and introduction of airborne contaminants. Salinity was maintained between 34 and 35 ppt by float switches which triggered pumps to transfer RODI freshwater housed in a 55-gal holding tank to the main tanks. An ultraviolet (UV) sterilizer (Coralife® Turbo Twist 9 W) controlled free-floating algae that could decrease light penetration to the corals.

Three light systems were used to provide PAR. The first was broad-spectrum (broad wavelength) high-intensity lighting derived from a 400 W metal halide ballast (PFO®) and two 10,000 K bulbs (Ushio®). Second, two 160 W actinic VHO bulbs (UVL lighting® T12) and a VHO (PFO®) ballast produced light in the 420 nm range to accommodate the energetic/photosynthetic needs of the zooxanthellae (Jeffrey and Haxo, 1968). Fans were installed to reduce excessive heat from the lights. Timers were used to simulate a natural 12:12 h light:dark photoperiod, with a duration of 5 h of high intensity metal halide lighting within the twelve-hour actinic lighting day. Two lunar lighting units (R2 Solutions® Moonlight 11000 MC) were also kept on continually.

2.6. Biotic nutrient controls

Live rock from Fiji was placed into the protein skimmers to serve as a biological filter, providing a high surface area for nitrifying bacteria. Algae associated with the live rock also helped to keep nitrogen (ammonia, nitrite, and nitrate) levels low. Turbo snails (*Turbo fluctuosa*) were introduced into the tanks to help prevent algal accumulation on the coral and tank. Because zooxanthellae photosynthate may be nutritionally limiting for corals (Johannes et al., 1970), supplemental food was provided to the corals to increase coral growth and survival (see Ferrier-Pagès et al., 2003; Lewis, 1974) three times per week, using Cyclop-Eeze (Liquid Life Argent®). The powerfilters (Aqua Clear®) were turned off during feeding for 30 min.

2.7. Data collection

Data were collected on the corals six times over a 115-day period. Measurements included total skeletal linear length (TSL; mm), living tissue linear length (LL; mm), weight (g), number of branches, and degree of polyp extension (extended vs. not extended). The threshold for counting new growth as a new branch was 5 mm. Three replicate

readings were taken of TSL (mm), LL (mm), and weight (g) during each sampling period to determine the precision of the measurement.

Wet weight was recorded for each coral fragment. The PVC support was dried and total weight of the block with coral was weighed on a digital Mettler scale. Polyp extension was measured three times per week, and recorded as fully extended, partially extended, or no extension.

After data collection, the positions of the PVC coral support structures were rotated in a uniform fashion within the tank to control for any potential position effects. During sampling, corals were photographed (front and back) for record-keeping purposes.

2.8. Length measurements

We made direct measurements of colony length. The variables were total skeletal length and living tissue length, measured using digital calipers.

2.9. Data analysis and graphics

Data were logged and collated using EXCEL 2003 and analyzed using SAS 2003 and BiomStat 3.3, using standard parametric *a priori* and *a posteriori/post-hoc* statistical tests (Sokal and Rohlf, 1981). Data were further analyzed and graphed using SigmaPlot 10.0. Diagrams were created using MS PowerPoint 2003 and Paint 2003.

3. Results

3.1. Control of experimental water quality

Mean phosphate levels in the tanks were 0.09 ± 0.018 , 0.20 ± 0.024 , and 0.50 ± 0.059 mg L⁻¹ (mean \pm 95% conf. intervals; target levels = 0.10, 0.20, and 0.50 mg L⁻¹; Fig. 4). Seawater temperature was maintained at 25 °C, pH 8.2, and 34–35‰ (Table 1). Calcium levels were maintained at 400 mg/L, not being permitted to fall below a mean level of 355–360 mg L⁻¹. (It is known that calcium levels > 360 mg L⁻¹ have no enhancing effects on coral calcification rates; Tambutté et al., 1996.) No ammonia was detected at any stage of the experiment, and nitrite and nitrate levels were negligible or absent during most of the experiment. Water quality parameters occasionally differed significantly between treatments (Table 1) but always remained well within the natural range known to support healthy coral growth (Kleypas et al., 1999).

3.2. Growth

Corals exposed to elevated phosphate levels exhibited increased growth rates. Corals maintained at the highest phosphate concentration (0.50 mg L⁻¹) had the highest total skeletal linear (TSL) growth rates (Fig. 5A). Total TSL growth at the highest phosphate concentration was 14.0 ± 6.20 mm over a period of 115 days, significantly greater than TSL growth at the lower phosphate levels (Fig. 5B). No difference was found in TSL between the 0.09 and 0.20 phosphate mg L⁻¹ treatments (3.9 ± 2.58 and 4.8 ± 5.35 mm, respectively). Between a point sixty-five days prior to initiation of the experiment and addition of phosphate to the tanks (t_0), fragments (ramets, or clones) of Colony A exhibited low or negligible TSL growth rates in all experimental tanks (Fig. 6). Once phosphate levels were increased, fragments exposed to the highest levels exhibited significantly greater growth rates.

Phosphate-enhanced growth through time, as measured by living length (LL) of coral tissue, was also significantly higher in the 0.50 mg L⁻¹ treatment than at lower concentrations (Fig. 7A). Total growth with respect to LL among the three treatments was 14.6 ± 5.43 mm of the highest phosphate concentration treatment (Fig. 7B) and was significantly greater than growth in the other




Fig. 4. Relationship between actual experimental concentrations of phosphate (0.09, 0.20, 0.50 mg L⁻¹) and target concentrations (0.10, 0.20, 0.50 mg L⁻¹). Mean shown with 95% confidence levels. Significant difference in phosphate concentrations between the 0.50 mg phosphate L⁻¹, 0.20 mg L⁻¹, and 0.09 mg L⁻¹ treatments ($p < 0.05$, one-way ANOVA; $p < 0.05$, Tukey's *post hoc* test; treatments that do not share a common letter are significantly different).

two treatments (4.6 ± 2.14 mm at 0.09 mg P L⁻¹, and 5.8 ± 4.80 mm at 0.20 mg P L⁻¹).

Weight of the corals increased significantly with time in all treatments (Fig. 8). Those corals subjected to the highest concentrations of phosphate (0.50 mg L⁻¹) accumulated weight at the highest rate. Weight accumulation was significantly different between all treatments.

Relative density (RD) was calculated according to the following formula:

$$RD = [(W_t/W_{max})/(TSL_t/TSL_{max})] - [(W_i/W_{max})/(TSL_i/TSL_{max})] + 1,$$

where RD = Relative Density, W = wet weight of individual fragment in g, TSL = total skeletal length in mm, i = initial measurement,

Table 1

Physico-chemical attributes of the experimental seawater. Mean \pm s.d. shown for pH, temperature (°C), salinity (‰), calcium (mg L⁻¹), ammonia-N (mg L⁻¹), nitrite (mg L⁻¹), and nitrate-N (mg L⁻¹). Calcium levels were returned to 400 mg L⁻¹ when required; mean weekly levels shown. Data analyzed by one-way ANOVA ($\alpha = 0.05$) followed by Tukey's *post hoc* test. Means within a category designated by a common letter are not significantly different ($p > 0.05$).

Variable	Phosphate level	Mean SD	N
pH	0.09 mg/L	8.22 ± 0.100^A	71
	0.20 mg/L	8.26 ± 0.094^B	71
	0.50 mg/L	8.25 ± 0.105^{AB}	71
Temperature	0.09 mg/L	25.35 ± 0.692^A	56
	0.20 mg/L	25.27 ± 0.586^A	56
	0.50 mg/L	25.30 ± 0.686^A	56
Salinity	0.09 mg/L	34.39 ± 1.073^A	56
	0.20 mg/L	34.57 ± 0.912^A	56
	0.50 mg/L	35.00 ± 0.638^B	55
Calcium	0.09 mg/L	360.88 ± 50.536^A	17
	0.20 mg/L	367.06 ± 32.262^A	17
	0.50 mg/L	375.29 ± 39.979^A	17
Ammonia	0.09 mg/L	$0.00 \pm N/A^A$	15
	0.20 mg/L	$0.00 \pm N/A^A$	14
	0.50 mg/L	$0.00 \pm N/A^A$	16
Nitrite	0.09 mg/L	0.03 ± 0.165^{AB}	45
	0.20 mg/L	$0.00 \pm N/A^A$	41
	0.50 mg/L	0.09 ± 0.212^B	44
Nitrate	0.09 mg/L	0.13 ± 0.168^A	14
	0.20 mg/L	0.11 ± 0.196^A	14
	0.50 mg/L	0.13 ± 0.198^A	14




Fig. 5. (A) Changes in total skeletal length (TSL; mm) in *Acropora muricata* under varying phosphate concentrations through time (days). Mean shown with 95% confidence levels. TSL growth is significantly higher at 0.50 mg phosphate L⁻¹ than in the 0.20 mg L⁻¹ and 0.09 mg L⁻¹ treatments ($p < 0.05$, repeated measures one-way nested ANOVA; treatments that do not share a common letter are significantly different). 0.09 mg L⁻¹ treatment: $p < 0.01$, linear regression, $Y = 0.028X + 0.185$, $r = 0.915$ (Pearson's product moment correlation coefficient, Sokal and Rohlf 1981); 0.20 mg L⁻¹ treatment: $p < 0.001$, $Y = 0.038X + 0.191$, $r = 0.985$; 0.50 mg L⁻¹ treatment: $p < 0.001$, $Y = 0.122X - 0.185$, $r = 0.997$. (B) Overall growth response indicated by total skeletal length (TSL₆ - TSL₀; mm) in *Acropora muricata* under different experimental concentrations. Mean shown with 95% confidence levels. TSL total growth gained is significantly higher at 0.50 mg phosphate L⁻¹ than in the 0.20 mg L⁻¹ and 0.09 mg L⁻¹ treatments ($p < 0.05$, one-way ANOVA; $p < 0.05$, Tukey's *post hoc* test).

t = measurement at time t , and \max = maximal observation for each variable per coral fragment. Relative density of corals from the highest phosphate concentration was found to be significantly lower than corals from the other two lower phosphate concentration treatments (Fig. 9).

There was no significant difference in production of new branches between the corals in the various phosphate enrichment treatments (1.1 ± 0.69 , branch production \pm 95% conf. limits at 0.09 mg L⁻¹ phosphate concentration; 1.3 ± 0.57 at 0.20 mg L⁻¹; 2.3 ± 2.37 at 0.50 mg L⁻¹).

No significant differences were detected in frequency of polyp extension among the experimental treatments (1.0 ± 0.01 - mean extension \pm 95% conf. limits - at 0.09 mg L⁻¹ phosphate concentration; 1.0 ± 0.01 at 0.20 mg L⁻¹; 1.0 ± 0.00 at 0.50 mg L⁻¹).

4. Discussion

There are sources of variance that we have eliminated here that are present in other studies, particularly field studies. Koop et al.'s (2001) experiment yielded some valuable results, but it was understood that experimental levels of nutrients added to the reef varied




Fig. 6. Changes in total skeletal length (TSL; mm) in *Acropora muricata* in Colony A under varying phosphate concentrations through time (days). Pre-experimental growth rates shown as well. Mean shown with 95% confidence levels. Colony A TSL growth is significantly higher at 0.50 mg phosphate L⁻¹ than the 0.20 mg L⁻¹ and 0.09 mg L⁻¹ treatments ($p < 0.05$, repeated measures one-way nested ANOVA; treatments that do not share a common letter are significantly different). 0.09 mg L⁻¹ treatment: $p < 0.05$, linear regression, $Y = 0.010X + 0.174$, $r = 0.658$ (Pearson's product moment correlation coefficient, Sokal and Rohlf 1981); 0.20 mg L⁻¹ treatment: $p < 0.001$, $Y = 0.014X + 0.837$, $r = 0.889$; 0.50 mg L⁻¹ treatment: $p < 0.001$, $Y = 1.018X + 2.365$, $r = 0.967$.




Fig. 7. (A) Changes in living tissue length (LL; mm) in *Acropora muricata* under varying phosphate concentrations through time (days). Mean shown with 95% confidence levels. LL growth is significantly higher at 0.50 mg phosphate L⁻¹ than in the 0.20 mg L⁻¹ and 0.09 mg L⁻¹ treatments ($p < 0.05$, repeated measures one-way nested ANOVA; treatments that do not share a common letter are significantly different). 0.09 mg L⁻¹ treatment: $p < 0.01$, linear regression, $Y = 0.038X - 0.445$, $r = 0.932$ (Pearson's product moment correlation coefficient, Sokal and Rohlf 1981); 0.20 mg L⁻¹ treatment: $p < 0.001$, $Y = 0.046X + 0.119$, $r = 0.982$; 0.50 mg L⁻¹ treatment: $p < 0.001$, $Y = 0.129X - 0.416$, $r = 0.999$. (B) Overall growth response in living tissue length through time (LL₆ - LL₀; mm) in *Acropora muricata* under different phosphate concentrations. Mean shown with 95% confidence levels. LL gained is significantly higher at 0.50 mg phosphate L⁻¹ than in the 0.20 mg L⁻¹ and 0.09 mg L⁻¹ treatments ($p < 0.05$, one-way ANOVA; $p < 0.05$, Tukey's *post hoc* test).




Fig. 8. Changes in weight (g) in *Acropora muricata* under varying phosphate concentrations through time (days). Mean shown with 95% confidence levels. Significant difference in weight-gain between the 0.50 mg phosphate L^{-1} , 0.20 mg L^{-1} , and 0.09 mg L^{-1} concentrations ($p < 0.05$, repeated measures one-way nested ANOVA). 0.09 mg L^{-1} treatment: $p < 0.001$, linear regression, $Y = 0.009X + 0.024$, $r = 0.994$, (Pearson's product moment correlation coefficient, Sokal and Rohlf 1981); 0.20 mg L^{-1} treatment: $p < 0.001$, $Y = 0.016X - 0.071$, $r = 0.983$; 0.50 mg L^{-1} treatment: $p < 0.001$, $Y = 0.024X - 0.149$, $r = 0.998$.

with the tide because of natural flushing. To avoid such problems, we took advantage of a controlled laboratory environment. Experimental levels of phosphate here were maintained very close to pre-defined target levels throughout the experiment through frequent monitoring, removing this potential source of variance. In addition, the coral colonies used here were determined to be genetically distinct, and showed no significant differences in growth rate prior to the addition of phosphate. This indicates that any treatment effects observed were not due to differences between the colonies. Thus, the differences in coral growth reported here are due primarily to differences in phosphate concentration and not to differences in other parameters such as genetics or aquaria conditions.

There was no difference in coral growth rates in the 0.09 and 0.20 mg L^{-1} phosphate concentration treatments, but at 0.50 mg L^{-1} , a significant difference did emerge. This indicates that a phosphate-




Fig. 9. Changes in relative density of *Acropora muricata* under varying phosphate concentrations through time (days). Relative density = $P_t [(W_i/W_{max})/(TSL_i/TSL_{max})] - P_i [(W_i/W_{max})/(TSL_i/TSL_{max})] + 1$, where i = initial, and t = measurement at time t . Mean shown with 95% confidence levels. Significant difference in relative density between the 0.50 mg phosphate L^{-1} vs. the 0.20 mg L^{-1} and 0.09 mg L^{-1} treatments ($p < 0.05$, repeated measures one-way nested ANOVA; treatments that share a common letter are not significantly different). 0.09 mg L^{-1} treatment: $p < 0.01$, linear regression, $Y = -0.0006X + 0.997$, $r = 0.958$, (Pearson's product moment correlation coefficient, Sokal and Rohlf 1981); 0.20 mg L^{-1} treatment: $p < 0.01$, $Y = -0.0008X + 0.982$, $r = 0.932$; 0.50 mg L^{-1} treatment: $p < 0.001$, $Y = -0.0034X + 0.973$, $r = 0.979$.

enhanced growth effect is manifested somewhere between 0.20 and 0.50 mg L^{-1} .

The graded response in weight accumulation which occurred in corals exposed to the experimental treatments indicates that, of the variables considered here, weight is the most sensitive to phosphate concentration. Our density measurements indicate that the rate of weight increase did not keep pace with the rate of skeletal extension, resulting in a more rapidly growing, but less dense coral skeleton at the highest phosphate concentration. The skeleton was becoming less dense, more porous, and most likely weaker. Most of the weight increase was, of course, due to accretion of calcium carbonate. A threshold effect was also detected in this case, occurring between 0.20 and 0.50 mg L^{-1} phosphate. Calcium carbonate deposition is clearly more tightly linked to ambient phosphate concentration than skeletal extension or live tissue length.

Acropora species appear to be tolerant of high phosphate levels and exhibit increased growth under these conditions, when compared to other corals (Bucher and Harrison, 2002; Koop et al., 2001; Steven and Broadbent, 1997). Bucher and Harrison (2002) suggest that *Acropora* growth may be less affected than other coral species when exposed to an increase in inorganic nutrient levels, in the absence of stresses such as reduced salinity, elevated sediments, or pollutants. Inter-specific variation in responses may also be due to different clades of *Symbiodinium* sp. Other potential factors include the presence (sometimes in high abundances) or absence of known or unknown enzymes (e.g., Ca^{2+} -ATPase, carbonic anhydrase).

4.1. Variation in results from earlier studies

Our study supports recent field studies (Bucher and Harrison, 2002; Koop et al., 2001; Steven and Broadbent, 1997) that demonstrated an increase in growth of *Acropora* species under conditions of increased phosphate levels, similar to those used here. On the other hand, Kinsey and Davies (1979) hypothesized that phosphate can inhibit coral growth and may cause a decrease in skeletal density.

In some earlier experiments, increased phosphate concentrations caused a reduction in coral calcification rates (Ferrier-Pagès et al., 2000), particularly in *A. cervicornis* (Renegar and Riegl, 2005). Variation in results may have been caused by the use of different experimental techniques, e.g., the use of small tanks (e.g. 5–8 L) with a high density of coral fragments. Here, we used large tanks (110 L) and a small number of small fragments in each, eliminating crowding and water quality problems. Adding food to eliminate any nutritional limitations (Johannes et al., 1970) also helped to support healthy coral growth.

Determining weight via a simple wet-weight technique has, we believe, benefited this study.

Past studies have often used the buoyant weight technique (Davies, 1989). Bucher et al. (1998), however, have questioned the accuracy of that technique due to variations in coral skeletal micro-density found between species and regions. In addition, it assumes that the micro-density of the coral skeleton is equivalent to that of pure aragonite (2.94 g/cm³). Significant variations in skeletal density have been found between *Acropora* species characterized by differences in chemistry or skeletal micro-architecture from different regions.

4.2. Mechanisms underlying effects of phosphate enrichment on coral growth

It is possible that the observed accelerated growth under phosphate-enriched conditions may be due to increased zooxanthellar populations and photosynthetic production within the coral. An increase in zooxanthellar populations within corals is known to occur under conditions of increased phosphate concentrations (Koop et al., 2001; Steven and Broadbent, 1997; Takabayashi, 1996). Phosphate is also known to increase chlorophyll levels (Bucher and Harrison, 2002; Koop et al., 2001) and photosynthetic rates (Ferrier-Pagès et al., 2000).

Increasing phosphate levels may have “fertilized” the zooxanthellae, stimulating their population growth. The stimulation of zooxanthellar cell production and coral growth by elevated phosphate levels has been reported by other investigators (Bucher and Harrison, 2002; Koop et al., 2001). As zooxanthellar production increases, the coral most likely receives additional translocation products such as photosynthetically derived ATP, which in turn allows for an increase in calcification and growth. Such an increase in ATP could enhance calcification by fueling active transcellular transport mechanisms (Al-Horani et al., 2003; Tambutté et al., 1996). An increase in ATP production could increase calcification by transporting additional calcium into the extracytoplasmic calcifying fluid (ECF) via the Ca^{2+} -ATPase-mediated $\text{Ca}^{2+}/2\text{H}^{+}$ exchange which occurs at the calciblastic epithelium (Fig. 10A). Therefore, an increase in calcium ions within the ECF could lead to increased calcification and growth rates.

An increase in the zooxanthellar population could disrupt the equilibrium of products exchanged between the host and symbiont. An influx of H^{+} ions produced by calcification can result in acidification of the environment. Furla et al. (1998) suggested that, under normal conditions, a stable intra-coelenteron pH is maintained by OH^{-} ions derived from photosynthesis. Tanaka et al. (2007) proposed that if photosynthesis increases, an alkalinizing effect could occur within the coelenteron, increasing calcification in order to generate additional H^{+} ions in order to neutralize elevated OH^{-} levels resulting in high growth rates. As a result, elevated levels of calcium ions may be transported to the calcification site.

The phosphate-enhanced coral growth we observed here may be due to an increase in active trans-cellular calcium transport resulting from expendable energy reserves, or calcification enhanced by alkalization within the coral (Fig. 10B).

4.3. Phosphate concentrations and skeletal density

Phosphate has been referred to as a “crystal poison” to the calcifying environment and the causative factor of coral density loss under nutrient enrichment (Kinsey and Davies, 1979; Risk and Sammarco, 1991; Simkiss, 1964). Our results support recent studies that have demonstrated an increase in porosity and a decrease in density under phosphate-enriched conditions (Koop et al., 2001). This condition may cause the coral to be vulnerable to breakage or more susceptible to natural wave action and colonization by internal bioeroders, further decreasing its skeletal integrity (Risk et al., 1995; Sammarco and Risk, 1990). A rapid “lean” linear growth and a trend towards the production of numerous branches would allow for increased surface area for photosynthesis and feeding. Although this trend would appear to be advantageous, as mentioned above, less heavily calcified corals may become brittle and more prone to breakage, and thus represent an unhealthy state.

We are suggesting that coral density may be decreased due to phosphate ions distorting the orderly array of ions on the crystal surface. Simkiss (1964) has proposed that phosphate may be binding at the site of calcification. Recent studies have confirmed this suggestion finding a correlation between phosphate within the coral skeleton and in the surrounding waters (Montagna et al. 2006). Raistrick's (1949) hypothesis suggested that each phosphate group is capable of replacing one carbonate ion in the crystal lattice (Simkiss, 1964). In this case, crystal distortion will occur because the foreign metaphosphate ions do not conform to the three-dimensional structure normally characteristic of the crystal lattice. As a result of this distortion, coral skeletal integrity may be compromised due to the production of a porous and structurally weaker calcium carbonate/calcium phosphate skeleton (Fig. 11). Although recent research suggests that this “anionic substitution” is responsible for only a small proportion of phosphate incorporated into the skeleton, the incorporation mechanism is still unclear and may vary between genera and forms of salt-water phosphorus (LaVigne et al., 2008; LaVigne et al., 2010).




Fig. 10. (A) Mechanism by which coral growth could be stimulated via phosphate-enhanced zooxanthellar production. Additional photosynthetically-derived ATP could allow for an increase of active calcium transport to the extracytoplasmic calcifying fluid (ECF), thereby stimulating increased calcification and growth rates (see Allemand et al. 2004; Furla et al. 2000; Gattuso et al. 1999; McConnaughey and Whelan 1997 for descriptions of the calcification process). (B) Mechanism by which phosphate-enhanced zooxanthellar population density can cause intra-coelenteric alkalization due to an increase in HCO_3^- derived photosynthesis. Corals maintain an optimum intra-coelenteron pH by increasing the influx of H^{+} ions produced by calcification. As a result of this, increased calcium ions are transported to the area of calcification, resulting in faster growth (see Tanaka et al. 2007; also see Weis et al. 1989; Furla et al. 1998).

To summarize, we suggest that a phosphate-induced increase in zooxanthellar populations may cause an increase in photosynthate and expendable energy reserves, and in turn permit increased calcification/growth. The increased photosynthesis would create a higher demand for carbonic anhydrase, thus increasing the pH in the intra-coelenteric space. This would also create a deficit of carbonate available for calcification which in turn would allow the phosphate ions to be used as a supplement to the carbonate ions while simultaneously helping to maintain an optimum pH. A lighter skeleton containing higher levels of calcium phosphate may then be formed, resulting in an increase in coral skeletal porosity.

4.4. Coral health in an eutrophic environment — experimental evidence for the Janus effect

Understanding the effects of nutrients on coral growth is important for maintaining the health of a coral reef ecosystem. It has been demonstrated here and elsewhere that corals can grow faster under high phosphate conditions (Bucher and Harrison, 2002; Koop et al., 2001; Steven and Broadbent, 1997). Higher coral growth rates were once considered to be a positive indicator of coral health (Neudecker, 1983; Rosen, 1981). Edinger et al. (2000) have suggested the opposite, however. Although individual corals on eutrophic reefs may grow faster, colonies are often susceptible to increased bioerosion pressure as well as competition for space with non-calcified organisms (also see Birkeland, 1987;

Edinger and Risk, 1994; Glynn, 1988; Hallock and Schlager, 1986; Risk and Sammarco, 1991; Risk et al., 1995; Sammarco and Risk, 1990). In fact, they claim that under nutrient enrichment conditions, coral growth rates and reef growth rates become decoupled. Edinger and Risk (1994; Edinger et al., 2000) have deemed this phenomenon “the Janus effect”, where an increase in coral growth rates cannot compensate for the additional pressures associated with eutrophic conditions affecting accretion rates, and reef growth is inhibited. In addition, coral skeleton integrity may be further compromised due to the production of a structurally inferior calcium phosphate skeleton. As a result, corals may be more prone to breakage resulting from human and storms. Exposed skeletons may allow for algal colonization and a potential shift in community structure.

In our study, the highest phosphate concentration produced no visible effects of stress on the corals, as indicated by polyp extension. In addition, no coral mortality was observed. Phosphate levels tested were much higher than those often found on natural reefs. This implies that increased phosphate alone may not lead to widespread reef mortality. The apparent health of the reef, however, may decline as conditions become more eutrophic reefs, resulting in high

bioerosion rates, high phytoplankton populations (Walker and Ormond, 1982), and increased benthic filamentous algal populations (Done, 1992). Thus, it would appear that reefs in nutrient-rich waters may survive the symptoms of global climate change, but their lack of resilience to mechanical stress may ultimately lead to disintegration – i.e., sea-level rise and increased storm frequencies and intensities, but only with difficulty, due to their weakened condition.

Acknowledgments

J.G. Dunn thanks Q. Fontenot, R. Boopathy, and A. Pierce for comments on the manuscript. R. Laird, L. Meng, S. Jackson, T. Widgeon, and E. Dunn contributed in many ways to the successful completion of the experimentation. J.G. Dunn also thanks his family for their inspiration and support. Many thanks to H. Harrison for constructing the diagrams (Figs. 10 and 11). This study was supported by the Coastal Restoration and Enhancement through Science and Technology (CREST) Program Fund, the Louisiana Office of Environmental Education, and the generous support of J. Brothers and B. Castagnos. [SS]




Fig. 11. Proposed mechanism by which an increase in coral growth and porosity may increase under conditions of increased phosphate, in a DIC-limited environment. (A) In an oligotrophic environment, calcium and carbonate bind to form a dense calcium carbonate skeleton. (B) Under a phosphate-enriched conditions, zooxanthellar populations increase, decreasing available carbonate available for coral calcification. Coral growth could continue if calcium binds with newly available phosphate, thereby complementing the depleted carbonate levels. Crystal distortion could occur since the foreign phosphate ions would not be expected to conform to the precise three-dimensional structural fit in the normal crystal lattice. As a result, a porous and structurally weaker calcium carbonate/calcium phosphate skeleton would be produced. Such a scenario would allow skeletal growth to continue, but with decreased density.

References

- Aeby, G.S., Bourne, D.G., Wilson, B., Work, T.M., 2011. Coral diversity and the severity of disease outbreaks: a cross-regional comparison of *Acropora* White Syndrome in a species-rich region (American Samoa) with a species-poor region (Northwestern Hawaiian Islands). *J. Mar. Biol.* doi:10.1155/2011/490198.
- Al-Horani, F.A., Al-Moghrabi, S.M., de Beer, D., 2003. The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*. *Mar. Biol.* 142, 419–426.
- Allemand, D., Ferrier-Pagès, C., Furla, P., Houlbrèque, F., Puverel, S., Reynaud, E., Tambutté, E., Tambutté, S., Zoccola, D., 2004. Biomineralisation in reef-building corals: From molecular mechanisms to environmental control. *C. R. Palevol* 3, 453–467.
- Aronson, R.B., Precht, W.F., 2001. White-band diseases and the changing face of Caribbean coral reef. *Hydrobiol* 460, 25–38.
- Atchison, A.D., 2005. Offshore oil and gas platforms as stepping stones for expansion of coral communities: A molecular genetic analysis. M.Sc. thesis, Louisiana State University, Baton Rouge, Louisiana.
- Atchison, A.D., Sammarco, P.W., Brazeau, D.A., 2008. Genetic connectivity in corals on the Flower Garden Banks and surrounding oil/gas platforms, Gulf of Mexico. *J. Exp. Mar. Biol. Ecol.* 365, 1–12.
- Baker, A.C., 2003. Flexibility and specificity in coral–algal symbiosis: Diversity, ecology, and biogeography of *Symbiodinium*. *Ann. Rev. Ecol. Syst.* 34, 661–689.
- Baker, A.C., Glynn, P.W., Riegl, B., 2008. Climate change and coral reef bleaching: An ecological assessment of long-term impacts, recovery trends, and future outlook. *Estuar. Coastal Shelf Sci.* 80, 435–471.
- Barnes, R.D., 1987. *Invertebrate Zoology*, 5th ed. Saunders College Publ, Philadelphia, PA.
- Barnes, R.S.K., Hughes, R.N., 1982. *An Introduction to Marine Ecology*. Blackwell Scientific Publications, Oxford.
- Bell, P.R., Lapointe, B.E., Elmetri, I., 2007. Reevaluation of ENCORE: Support for the eutrophication threshold model for coral reefs. *Ambio* 36, 416–424.
- Birkeland, C.E., 1987. Nutrient availability as a major determining factor in differences among coastal hard-substratum communities in different regions of the tropics. In: Birkeland, C.E. (Ed.), *Comparison between Atlantic and Pacific Tropical Marine Coastal Ecosystems: Community Structure, Ecological Processes, and Productivity*. UNESCO/COMAR.
- Borneman, E.H., 2001. *Aquarium Corals: Selection, Husbandry, and Natural History*. T.F.H. Publ., Inc, Neptune City, New Jersey.
- Brazeau, D.A., Sammarco, P.W., Gleason, D.F., 2005. A multi-locus genetic assignment technique to assess sources of *Agaricia agaricites* larvae on coral reefs. *Mar. Biol.* 147, 1141–1148.
- Bruno, J.F., Petes, L.E., Harvell, C.D., Hettinger, A., 2003. Nutrient enrichment can increase the severity of coral diseases. *Ecol. Lett.* 6, 1056–1061.
- Bucher, D.J., Harrison, P.L., 2002. Growth response of the reef coral *Acropora longicyathus* to elevated inorganic nutrients: Do responses to nutrients vary among coral taxa? *Proc. 9th Int. Coral Reef Symp.* Vol. 1, pp. 443–448. Bali, Indonesia.
- Bucher, D.J., Harriott, V.J., Roberts, L.G., 1998. Skeletal micro-density, porosity and bulk density of acroporid corals. *J. Exp. Mar. Biol. Ecol.* 228, 117–136.
- Costa Jr., O.S., Leão, Z.M.A.N., Nimmo, M., Attrill, M.J., 2000. Nutritification impacts on coral reefs from northern Bahia, Brazil. *Hydrobiol.* 440, 307–315.
- Cruz-Piñón, G., Carriart-Ganivet, G.P., Espinoza-Avalos, J., 2003. Monthly skeletal extension rates of the hermatypic corals *Montastraea annularis* and *Montastraea faveolata*: Biological and environmental controls. *Mar. Biol.* 143, 491–500.
- Davies, P.S., 1989. Short-term growth measurements of corals using an accurate buoyant weighing technique. *Mar. Biol.* 101, 389–395.
- DeGeorges, A., Goreau, T.J., Reilly, B., 2010. Land-sourced pollution with an emphasis on domestic sewage: Lessons from the Caribbean and implications for coastal development on Indian Ocean and Pacific coral reefs. *Sustainability* 2, 2919–2949.
- Done, T.J., 1992. Phase shifts in coral reef communities and their ecological significance. *Hydrobiol* 247, 121–132.
- Edinger, E.N., Risk, M.J., 1994. Oligocene–Miocene extinction and geographic restriction of Caribbean corals: Roles of turbidity, temperature, and nutrients. *Palaios* 9, 576–598.
- Edinger, E.N., Limmon, G.V., Jompa, J., Widjtmoko, W., Heikoop, J.M., Risk, M.J., 2000. Normal coral growth rates on dying reefs: Are coral growth rates good indicators of reef health? *Mar. Pollut. Bull.* 5, 404–425.
- Fabrizius, K.E., 2005. Effects of terrestrial runoff on the ecology of corals and coral reefs: Review and synthesis. *Mar. Pollut. Bull.* 50, 125–146.
- Falkowski, P.G., Dubinsky, Z., Muscatine, L., Porter, J.W., 1984. Light and the bioenergetics of a symbiotic coral. *Biosci* 34, 705–709.
- Ferrier-Pagès, C., Gattuso, J.-P., Dallot, S., Jaubert, J., 2000. Effect of nutrient enrichment on growth and photosynthesis of the zooxanthellate coral *Stylophora pistillata*. *Coral Reefs* 19, 103–113.
- Ferrier-Pagès, C., Witting, J., Tambutté, E., Sebans, K.P., 2003. Effect of natural zooplankton feeding on the tissue and skeletal growth of the scleractinian coral *Stylophora pistillata*. *Coral Reefs* 22, 229–240.
- Finkel, C.W., Charlier, R.H., 2003. Sustainability of subtropical coastal zones in southeastern Florida: Challenges for urbanized coastal environments threatened by development, pollution, water supply, and storm hazards. *J. Coast. Res.* 19, 934–943.
- Fishelson, L., 1973. Ecology of coral reefs in the Gulf of Aqaba (Red Sea) influenced by pollution. *Oecologia* 12, 55–67.
- Furla, P., Benazet-Tambutté, S., Jaubert, J., Allemand, D., 1998. Functional polarity of the tentacle of the sea anemone *Anemonia viridis*: Role in inorganic carbon acquisition. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 274, 303–310.
- Furla, P., Galgani, I., Durand, I., Allemand, D., 2000. Sources and mechanisms of inorganic carbon transport for coral calcification and photosynthesis. *J. Exp. Biol.* 203, 3445–3457.
- Furla, P., Richier, S., Allemand, D., 2011. In: Dubinsky, Z., Stambler, N. (Eds.), *Physiological Adaptation to Symbiosis in Cnidarians: Coral Reefs: An Ecosystem in Transition*. Springer, New York, pp. 187–195.
- Futuyama, D.J., 1998. *Evolutionary Biology*, 3rd Ed. Sinauer Associates, Sunderland, Mass.
- Gattuso, J.-P., Allemand, D., Frankignoulle, M., 1999. Photosynthesis and calcification at cellular, organismal, and community levels in coral reefs: A review on interactions and control by carbonate chemistry. *Am. Zool.* 39, 160–183.
- Glynn, P., 1988. El Niño–Southern Oscillation 1982–1983: Near-shore population, community, and ecosystem responses. *Ann. Rev. Ecol. Syst.* 19, 309–345.
- Green, E.P., Bruckner, A.W., 2000. The significance of coral disease epizootiology for coral reef conservation. *Biol. Conserv.* 96, 347–361.
- Guzner, B., Novplansky, A., Shalit, O., Chadwick, N.E., 2010. Indirect impacts of recreational scuba diving: Patterns of growth and predation in branching stony corals. *Bull. Mar. Sci.* 86, 727–742.
- Hallock, P., Schlager, W., 1986. Nutrient excess and the demise of coral reefs and carbonate plateforms. *PALAIOS* 1, 389–398.
- Harrison, P.L., Ward, S., 2001. Elevated levels of nitrogen and phosphorus reduce fertilization success of gametes from scleractinian reef corals. *Mar. Biol.* 139, 1057–1068.
- Hawkins, J.P., Roberts, C.M., 1994. The growth of coastal tourism in the Red Sea: Present and future effects on coral reefs. *Ambio* 23, 503–508.
- Hay, M.E., Rasher, D.B., 2010. Coral reefs in crisis: Reversing the biotic death spiral. *Biol. Rep.* doi:10.3410/B2-71 F1000.
- Hoegh-Guldberg, O., 1999. Climate change, coral bleaching, and the future of the world's coral reefs. *Mar. Freshw. Res.* 50, 839–866.
- Hughes, T.P., 1994. Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science* 265, 1547–1551.
- Hughes, T.P., Connell, J.H., 1999. Multiple stressors on coral reefs: A long-term perspective. *Limnol. Oceanogr.* 44, 932–940.
- Hughes, T.P., Graham, N.A.J., Jackson, J.B.C., Mumby, P.J., Steneck, R.S., 2010. Rising to the challenge of sustaining coral reef resilience. *Trends Ecol. Evol.* 25, 633–642.
- Hunter, C.L., Evans, C.W., 1995. Coral reefs in Kaneohe Bay, Hawaii: Two centuries of western influence and two decades of data. *Bull. Mar. Sci.* 57, 501–515.
- Hunter, C.L., Stephenson, M.D., 1999. Pollution impacts on biodiversity in coral reef ecosystems of central Pacific islands. In: Eldredge, L.G., Maragos, J.E., Holthus, P.F., Takeuchi, H.F. (Eds.), *Marine and Coastal Biodiversity in the Tropical Island Pacific Region*. Pac. Sci. Assn, Honolulu, HI, pp. 419–440.
- Jackson, J.B.C., 1991. Adaptation and diversity of reef corals: Patterns result from species differences in resource use and life histories and from disturbances. *Biosci* 41, 475–482.
- Jeffrey, S.W., Haxo, F.T., 1968. Photosynthetic pigments of symbiotic dinoflagellates (zooxanthellae) from corals and clams. *Biol. Bull.* 135, 149–165.
- Jinendradasa, S.S., Ekaratne, S.U.K., 2002. Linear Extension of *Acropora muricata* (Dana) at Selected Reef Locations in Sri Lanka: *Proc. 9th Int. Coral Reef Symp.*, Vol. 1, pp. 417–420. Bali, Indonesia.
- Johannes, R.E., Coles, S.L., Kuenzel, N.T., 1970. The role of zooplankton in the nutrition of some scleractinian corals. *Limnol. Oceanogr.* 15, 579–586.
- Juhasz, A., Ho, E., Bender, E., Fong, P., 2010. Does use of tropical beaches by tourists and island residents result in damage to fringing coral reefs? A case study in Moorea French Polynesia. *Mar. Pollut. Bull.* 60, 2251–2256.
- Kinsey, D.W., Davies, P.J., 1979. Effects of elevated nitrogen and phosphorus on coral reef growth. *Limnol. Oceanogr.* 24, 935–940.
- Kleypas, J.A., McManus, J.W., Meñez, L.A.B., 1999. Environmental limits to coral reef development: Where do we draw the line? *Am. Zool.* 39, 146–159.
- Knowlton, N., Jackson, J.B.C., 2008. Shifting baselines, local impacts, and global change on coral reefs. *Plos Biol.* 6, e54. doi:10.1371/journal.pbio.0060054.
- Koop, K., Booth, D., Broadbent, A., Brodie, J., Bucher, D., Capone, D., Coll, J., Dennison, W., Erdmann, M., Harrison, P., Hoegh-Guldberg, O., Hutchings, P., Jones, G.B., Larkum, A.W.D., O'Neal, J., Steven, A., Tentori, E., Ward, S., Williamson, J., Yellowlees, D., 2001. ENCORE: the effect of nutrient enrichment on coral reefs. Synthesis of results and conclusions. *Mar. Pollut. Bull.* 42, 91–120.
- Lapointe, B.E., O'Connell, J.D., Garrett, G.S., 1990. Nutrient couplings between on-site sewage disposal systems, groundwaters, and nearshore surface waters of the Florida Keys. *Biogeochem* 10, 289–307.
- Lapointe, B.E., Langton, R., Day, O., Potts, A.C., 2003. Integrated water quality and coral reef monitoring on fringing reefs of Tobago: Chemical and ecological evidence of sewage-driven eutrophication in the Buccoo reef complex. *Proc. Gulf Caribb. Fish. Inst.* 54, 457–472.
- Lapointe, B.E., Barile, P.J., Matzie, W.R., 2004. Anthropogenic nutrient enrichment of seagrass and coral reef communities in the lower Florida Keys: Discrimination of local versus regional nitrogen sources. *J. Exp. Mar. Biol. Ecol.* 308, 23–58.
- Lapointe, B.E., Langton, R., Bedford, B.J., Potts, A.C., Day, O., Hu, C., 2010. Land-based nutrient enrichment of the Buccoo reef complex and fringing coral reefs of Tobago, West Indies. *Mar. Pollut. Bull.* 60, 334–343.
- Laurance, W.F., 2010. Habitat destruction: Death by a thousand cuts. In: Navjot, S., Sodhi, S., Ehrlich, P.R. (Eds.), *Conservation Biology for All*. Oxford Univ Press, New York, pp. 73–87.
- LaVigne, M., Field, M.P., Angnostou, E., Grottolli, A., Wellington, G., Sherrell, R., 2008. Skeletal P/Ca tracks upwelling in Gulf of Panama coral: Evidence for a new seawater phosphate proxy. *Geophys. Res. Lett.* 35, L05604.
- LaVigne, M., Matthews, K.A., Grottolli, A.G., Cobb, K.M., Anagnostou, E., Cabioch, G., Sherrell, R.M., 2010. Coral skeleton P/Ca proxy for seawater phosphate: Multi-colony calibration with a contemporaneous seawater phosphate record. *Geochim. Cosmochim. Acta* 74, 1282–1293.
- Lewis, J.B., 1974. The importance of light and food upon the early growth of the reef coral *Favia fragum* (Esper). *J. Exp. Mar. Biol. Ecol.* 15, 299–304.

- McConnaughey, T.A., Whelan, J.F., 1997. Calcification generates protons for nutrient and bicarbonate uptake. *Earth Sci. Rev.* 42, 95–117.
- McManus, J.W., 1997. Tropical marine fisheries and the future of coral reefs: A brief review with emphasis on Southeast Asia. *Coral Reefs* 16, 121–127.
- Miller, G.T., 1999. *Living in the Environment: Principles, Connections, and Solutions*. Brooks/Cole Publishing Company, Pacific Grove.
- Moberg, F., Folke, C., 1999. Ecological goods and services of coral reef ecosystems. *Ecol. Econ.* 29, 215–233.
- Montagna, P., McCulloch, M., Taviani, M., Mazzoli, C., Vendrell, B., 2006. Phosphorus in cold-water corals as a proxy for seawater nutrient chemistry. *Sci* 312, 1788–1791.
- Muscantine, L., 1990. The role of symbiotic algae in carbon and energy flux in reef corals. *Ecosystems of the World 25: Coral Reefs*. Elsevier, Amsterdam.
- Muscantine, L., Cernichari, E., 1969. Assimilation of photosynthetic products of zooxanthellae by a reef coral. *Biol. Bull.* 137, 506–523.
- Muscantine, L., McCloskey, L.R., Marian, R.E., 1981. Estimating the daily contribution of carbon from zooxanthellae to coral animal respiration. *Limnol. Oceanogr.* 26, 601–611.
- Muscantine, L., Falkowski, P.G., Dubinsky, Z., Cook, P.A., McCloskey, L.R., 1989. The effect of external nutrient resources on the population dynamics of zooxanthellae in a reef coral. *Proc. R. Soc. Lond. Ser. B. Biol. Sci.* 236, 311–324.
- Neudecker, S., 1983. Growth and survival of scleractinian corals exposed to thermal effluents in Guam: Proc. 4th Int. Coral Reef Symp., Vol. 1, pp. 173–180. Manila, Philippines.
- Patterson, K.L., Porter, J.W., Ritchie, K.B., Polson, S.W., Mueller, E., Peters, E.C., Santavy, D.L., Smith, G.W., 2002. The etiology of white pox, a lethal disease of the Caribbean elkhorn coral, *Acropora palmata*. *Proc. Natl. Acad. Sci.* 99, 8725–8730.
- Porter, J.W., 1976. Autotrophy, heterotrophy, and resource partitioning in Caribbean reef-building corals. *Am. Nat.* 110, 731–742.
- Raistrick, B., 1949. The influence of foreign ions on crystal growth from solution. 1. The stabilization of the super-saturation of calcium carbonate solutions by anions possessing O–P–O–P–O chains. *Discip. Faraday Soc.* 5, 234–237.
- Rasmussen, C.E., 1986. An investigation of morphological changes, Low Isles, Northern Great Barrier Reef, Australia. Unpublished Honours Thesis, Dept. Geography, James Cook University NQ, Townsville, Qld., Australia.
- Rasmussen, C.E., 1988. Effects Carried by Mainland Runoff on Reefs of the Cairns Area: A Research Plan and Preliminary Results. Tech. Report. Great Barrier Marine Park Authority, Townsville, Qld., Australia, pp. 66–92.
- Rasmussen, C., Ness, S., Cuff, C., 1993. In: Richmond, R.H. (Ed.), A correlation between the El Niño–Southern oscillation and crystallographic variability in corals of the Great Barrier Reef, Australia. *Proc. 7th Int. Coral Reef Symp.*, Vol. 1. Univ. Guam Press, UOG Station, Guam, pp. 211–220.
- Reaka-Kudla, M.L., 1997. *Understanding and Protecting our Biological Resources*. Biodiversity II. Joseph Henry Press, Wash., DC.
- Renegar, D.A., Riegl, B.M., 2005. Effect of nutrient enrichment and elevated CO₂ partial pressure on growth rate of Atlantic scleractinian coral *Acropora cervicornis*. *Mar. Ecol. Prog. Ser.* 293, 69–76.
- Reopanichkul, P., Carter, R.W., Worachananant, S., Crossland, C.J., 2010. Wastewater discharge degrades coastal waters and reef communities in southern Thailand. *Mar. Environ. Res.* 69, 287–296.
- Risk, M.J., Sammarco, P.W., 1991. Cross-shelf trends in skeletal density of the massive coral *Porites lobata* from the Great Barrier Reef. *Mar. Ecol. Prog. Ser.* 69, 195–200.
- Risk, M.J., Sammarco, P.W., Edinger, E.N., 1995. Bioerosion in *Acropora* across the continental shelf of the Great Barrier Reef. *Coral Reefs* 14, 79–86.
- Roberts, C.M., McClean, C.J., Veron, J.E.N., Hawkins, J.P., Allen, G.R., McAllister, D.E., Mittermeier, C.G., Schueler, F.W., Spalding, M., Wells, F., Vynne, C., Werner, T.B., 2002. Marine biodiversity hotspots and conservation priorities for tropical reefs. *Sci* 295, 1280–1284.
- Rosen, B.R., 1981. The Tropical High Diversity Enigma – The Coral's Eye View. *Chance, Change, and Challenge: The Evolving Biosphere*. Brit. Mus. (Nat. Hist.), Cambridge Univ. Press, Cambridge, London.
- Rothenberger, P., Blondeau, J., Cox, C., Curtis, S., Fisher, W., et al., 2008. The state of coral reef ecosystems of the U.S. Virgin Islands. In: Waddell, J.E., Clarke, A.M. (Eds.), *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2008*. : NOAA Tech. Mem., 73. NOS NCCOS, pp. 29–73.
- Sammarco, P.W., 1980. *Diadema* and its relationship to coral spat mortality: grazing, competition, and biological disturbance. *J. Exp. Mar. Biol. Ecol.* 45, 245–272.
- Sammarco, P.W., 1982a. Echinoid grazing as a structuring force in coral reef communities: whole reef manipulations. *J. Exp. Mar. Biol. Ecol.* 61, 31–55.
- Sammarco, P.W., 1982b. Effects of grazing by *Diadema antillarum* Philippi (Echinodermata: Echinoidea) on algal diversity and community structure. *J. Exp. Mar. Biol. Ecol.* 65, 83–105.
- Sammarco, P.W., Risk, M.J., 1990. Large-scale patterns in internal bioerosion of *Porites*: cross continental shelf trends on the Great Barrier Reef. *Mar. Ecol. Prog. Ser.* 59, 145–156.
- Schutte, V.G.W., Selig, W.R., Bruno, J.F., 2010. Regional spatio-temporal trends in Caribbean coral reef benthic communities. *Mar. Ecol. Prog. Ser.* 402, 115–122.
- Sebens, K.P., 1994. Biodiversity of coral reefs: What are we losing and why? *Am. Zool.* 34, 115–133.
- Simkiss, K., 1964. Phosphates as crystal poisons of calcification. *Biol. Rev.* 39, 487–504.
- Smith, S.V., Kimmerer, W.J., Laws, E.A., Brock, R.E., Walsh, T.W., 1981. Kaneohe Bay sewage diversion experiment: Perspectives on ecosystem responses to nutritional perturbation. *Pac. Sci.* 35, 279–395.
- Sokal, R.R., Rohlf, F.J., 1981. *Biometry*, 2nd Ed. W. H. Freeman, New York.
- Stambler, N., 2011. Marine microalgae/cyanobacteria-invertebrate symbiosis. In: Seckbach, J., Dubinsky, Z. (Eds.), *All Flesh is Grass: Cellular Origin, Life in Extreme Habitats, and Astrobiology* 16, Pt. 6. Springer, New York, pp. 383–414.
- Stambler, N., Popper, N., Dubinsky, Z., Stimson, J., 1991. Effects of nutrient enrichment and water motion on the coral *Pocillopora damicornis*. *Pac. Sci.* 45, 299–307.
- Steven, A.D.L., Broadbent, A.D., 1997. Growth and metabolic responses of *Acropora palifera* to long-term nutrient enrichment: Proc. 8th Int. Coral Reef Symp., Vol. 1, pp. 867–872. Panama.
- Suzuki, A., Nakamori, T., Kayanne, H., 1995. The mechanism of production enhancement in coral reef carbonate systems: Model and empirical results. *Sed. Geol.* 99, 259–280.
- Takabayashi, M., 1996. Impacts of in situ inorganic enrichment on the scleractinian corals, *Stylophora pistillata* and *Pocillopora damicornis*. M.Sc. thesis, Univ. Sydney, Sydney, Australia.
- Tambutté, E., Allemand, D., Mueller, E., Jaubert, J., 1996. A compartmental approach to the mechanism of calcification in hermatypic corals. *J. Exp. Biol.* 199, 1029–1041.
- Tameler, J., Rajasuriya, A., 2008. Status of coral reefs in South Asia: Bangladesh Chago, India, Maldives, and Sri Lanka. In: Wilkinson, C. (Ed.), *Status of Coral Reefs of the World*, pp. 119–130. 2008.
- Tanaka, Y., Miyajima, T., Koike, I., Hayashibara, T., Ogawa, H., 2007. Imbalanced coral growth between organic tissue and carbonate skeleton caused by nutrient enrichment. *Limnol. Oceanogr.* 52, 1139–1146.
- Trench, R.K., 1993. Microalgal-invertebrate symbiosis: A review. *Endocytobiol. Cell Res.* 9, 135–175.
- Walker, D.I., Ormond, R.F.G., 1982. Coral death from sewage and phosphate pollution at Aqaba, Red Sea. *Mar. Pollut. Bull.* 13, 21–25.
- Wallace, C.C., 1999. *Staghorn Corals of the World: A Revision of the Genus Acropora*. CSIRO Publishing, Collingwood, Victoria, Australia. 438 p.
- Weis, V.M., Smith, G.J., Muscatine, L., 1989. A “CO₂ supply” mechanism in zooxanthellate cnidarians: Role of carbonic anhydrase. *Mar. Biol.* 40, 598–605.
- Wild, C., Hoegh-Guldberg, O., Naumann, M.S., Colombo-Pallotta, M.F., Ateweberhan, M., Fitt, W.K., Iglesias-Preto, R., Palmer, C., Bythell, J.C., Ortiz, J.-C., Loya, Y., van Woesik, R., 2010. Climate change impedes scleractinian corals as primary reef ecosystem engineers. *Mar. Freshw. Res.* 62, 205–215.
- Williams, D.E., Miller, M.W., 2005. Coral disease outbreak: Pattern, prevalence, and transmission in *Acropora cervicornis*. *Mar. Ecol. Prog. Ser.* 301, 119–128.
- Wood, E.M., 2001. Collection of Coral Reef Fish for Aquaria: Global Trade, Conservation Issues, and Management Strategies: *Mar. Conserv. Soc.*, UK. 80 pp.
- Wright, O.P., Marshall, A.T., 1991. Calcium transport across the isolated oral epithelium of scleractinian corals. *Coral Reefs* 10, 37–40.
- Yonge, C.M., Yonge, M.J., Nicholls, A.G., 1932. Studies on the physiology of corals. VI. The relationship between respiration in corals and the production of oxygen by their zooxanthellae. *Sci. Rep. Great Barr. Reef Exped.* 1, 213–251.
- Zakai, D., Chadwick-Furman, N.E., 2002. Impacts of intensive recreational diving on reef corals at Eilat, northern Red Sea. *Biol. Conserv.* 105, 179–187.